

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

EQUIPMENT MAINTENANCE AND USE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy water or 1% solution of ammonia, followed by several clear water rinses.

COMPATIBILITY

Compatible with biological control agents and can be used in Integrated Pest Management Programs (IPM).

RESISTANT WEEDS WARNING

Agricrop AFFRAY 300 Herbicide (AFFRAY 300) is a member of the phenoxys group of herbicides. Agricrop AFFRAY 300 has the disruptors of plant cell growth mode action. For weed resistance management AFFRAY 300 is a Group I herbicide.

Some naturally occurring weed biotypes resistant to Agricrop AFFRAY 300 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Agricrop AFFRAY 300 or other disruptors of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Agricrop Pty Ltd accepts no liability for any losses that may result from the failure of Agricrop AFFRAY 300 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds.

DO NOT spray cereals if lucerne is present.

DO NOT spray crops or weeds outside the stages indicated in 'Critical Comments' as damage, loss of yield or inadequate weed control may result.

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops, such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (e.g. Kurrajongs, Belahs and Eucalypts).

Avoid use on Sown Pastures.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 131126).

MSDS

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

Agricrop Pty. Ltd. ('Agricrop') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Agricrop's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Agricrop has any authority to add to or alter these conditions.

NRA Approval No.: 56510/1002

In case of Emergency Phone 1800 033 111 All hours Australia wide

AGRICROP PTY LTD A.C.N. 100 473 309

1569 Wynnum Rd TINGALPA QLD 4173

Tel (07) 3348 4113 Fax (07) 3348 2702



BATCH NO.:
DATE OF MANUFACTURE:

CAUTION

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

AGRICROP AFFRAY 300 HERBICIDE

ACTIVE CONSTITUENT: 300g/L 2,4-D
SOLVENT: 259g/L HYDROCARBON LIQUID

GROUP I HERBICIDE

NET CONTENTS

20L

For the integrated control of Groundsel Bush, Mother-of-Millions, Noogoora Burr, Bathurst Burr, Water Hyacinth and other weeds as listed in the Directions for Use table.

AGRICROP PTY LTD A.C.N. 100 473 309

1569 Wynnum Road TINGALPA QLD 4173

Tel (07) 3348 4113 Fax (07) 3348 2702

DIRECTIONS FOR USE

RESTRAINTS: Avoid use in sown pasture

Situation	Weeds Controlled	State	Rate	Critical Comments
Non-crop & Pasture	Groundsel Bush	Qld & NSW only	10L/ha	Helicopter application. Raindrop D8 nozzles angled back at 45° to minimise spray drift. Apply in no less than 100L/ha.
			100mL/10L water	Knapsack spray. Thorough even coverage of the plant is necessary.
			1L/10L water	Sprinkler sprayer. Coverage: 1L spray solution/100m².
	33mL/L kerosene or mineral turpentine		Basal bark spray: Apply the mixture around the base of the stem to a height of 25cm above the ground. Make sure all the stem to this height is well wetted. Cut stump: Cut the stem as close to the ground as possible and thoroughly wet the fresh cut stem with the mixture.	
	Mother-of-Millions		70mL/10L water	Knapsack spray. Thorough even coverage of leaves and plantlets is necessary.
			7L/1000L water/ha	Hand Gun. Thorough even coverage of leaves and plantlets is necessary.
	Noogoora Burr Bathurst Burr		3.3L/ha	Helicopter application or sprinkler sprayer.
	Castor Oil Plant	Qld. only	1L/10L kerosene	Apply as a basal bark treatment.
	Galvanised Burr Chinese Celtis (Elm) seedlings		1L/100L water	Apply as an overall spray on young plants when actively growing.
	Wild Tobacco Tree		1L/40L kerosene	Apply as a cut stump or basal bark when plant actively growing.
	Balloon Cotton Datura (seedlings) Sida Retusa Creeping Lantana Ageratum spp. Fireweed		7L/1000L water	Apply as an overall spray when plants actively growing.
	Saffron Thistle		1L/100L water	Apply as an overall spray when plants actively growing.
	Waterways, non-potable water, drains, dam margins, lakes & streams		NSW, Qld, WA and NT only	50mL/10L water
			1L/200L water	Hand Gun: Coverage: 200L spray solution/1000m². Avoid causing submersion of sprayed plants.
			1L/20L water	Sprinkler sprayer. Coverage: 20L spray solution/1000m².
			5L/200L water/ha	Helicopter application. Raindrop D8 nozzles angled back at 45° to minimise spray drift.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

AGRICROP

AGRICROP